

Name: Katsumi Miyauchi, M.D. PhD

Present Position:

Professor, Department of Cardiovascular Medicine
Juntendo Tokyo Geriatric Medical Center
Associate Director, Juntendo Research Support Center
Juntendo University, School of medicine

Office Address:

Cardiovascular Medicine ; Juntendo University Hospital
2-1-1 Hongo, Bunkyo-ku, Tokyo, Japan
Phone: 81-3-5802-1056; Fax: 81-3-5689-0627)

Education:

M.D.- Nagasaki University.— 1978-1984,
PhD-Juntendo University-1998-

Residency,

3rd Department of Internal Medicine Nagasaki University. —1984-1986

Fellowship

Department of Cardiology Juntendo University 1986-1988

1997:

Chief in Catheter laboratory

1998-2000:

Mayo Clinic Research Fellow. Rochester, Minnesota, USA

2001-2015:

Chief in Catheter laboratory in Juntendo University

2008:

Associate Professor Cardiovascular Medicine, Juntendo University

2016:

Professor, Department of Cardiovascular Medicine, Juntendo Tokyo Geriatric
Medical Center

Editor Board

Japanese Association of Cardiovascular Intervention and Therapeutics (CVIT)
Japanese College of Cardiology

Council Board of Professional Societies

Japanese Circulation Society
Japanese College of Cardiology
Japanese Association of Cardiovascular Intervention and Therapeutics
Japanese Society of Cardiovascular Imaging and Dynamics

Membership in Professional Societies:

Japanese Society of Internal Medicine, Japanese Circulation Society, Japanese
College of Cardiology, Japan Atherosclerosis Society, Japanese Coronary
Association, Japanese Society of Cardiovascular Imaging and Dynamics

Awards 2003 7th Cardiovascular Radiation Therapy (CRT), Washington DC Best Abstract Award
Efficacy of Activation of Peroxisome Proliferative-Activated Receptors (PPAR)- on Inflammatory
Response and Neointimal Thickening After Coronary Angioplasty

Licenser and Board Certification:

Japanese National Board of Medicine

Japanese Board of Cardiology

1. Kawano H, Miyauchi K, Okada R, Daida H, Yokoi H, Miyano H, Takaya J, Satoh H, Yamaguchi H, Suda K: Histopathological study of cardiac rupture following myocardial infarction with and without thrombolytic therapy. *J Cardiol.* 1994; 24: 249-255 (2.566)
2. Miyauchi K, Kawai S, Okada R, Yamaguchi H: Limitations of angiotensin-converting enzyme inhibitor in restenosis of a deep arterial injury model. *Jpn Circ J.* 1998; 62: 53-60, (3.685)
3. Miyauchi K, Aikawa M, Tani T, NakAmerican Heart Associatiiora K, Kawai S, Nagai R, Okada R, Yamaguchi H: Effects of probucol on smooth muscle cell proliferation and dedifferentiation after vascular injury in rabbits: possible role of PDGF. *Cardiovasc Drugs Ther.* 1998; 12: 251-260 (2.952)
4. Kimura T, Miyauchi K, Yamagami S, Daida H, Yamaguchi H: Local delivery infusion pressure is a key determinant of vascular damage and intimal thickening. *Jpn Circ J.* 1998; 62: 299-304 (3.685)
5. Kwon HM, Sangiorgi G, Spagnoli LG, Miyauchi K, Holmes DR Jr, Schwartz RS, Lerman A: Experimental hypercholesterolemia induces ultrastructural changes in the internal elastic lamina of porcine coronary arteries. *Atherosclerosis.* 1998 ; 139: 283-289 (3.971)
6. Yamagami S, Miyauchi K, Kimura T, Goh Y, Daida H, Yamaguchi H: Effects of the thromboxane A2 receptor antagonist on platelet deposition and intimal hyperplasia after balloon injury. *Jpn Heart J.* 1999; 40: 209-213 (3.685)
7. Kantor B, McKenna CJ, Cacchitolo JA, Miyauchi K, Reeder GS, Mullany CJ, Schaff HV, Holmes DR Jr, Schwartz RS: Transmyocardial and percutaneous myocardial revascularization: current and future role in the treatment of coronary artery disease. *Mayo Clin Proc.* 1999; 74: 585-592 (5.812)
8. Hamasaki S, Suwadi J, Higano ST, Miyauchi K, Holmes DR Jr, Lerman A: Attenuated coronary flow reserve and vascular remodeling in patients with hypertension and left ventricular hypertrophy. *J Am Coll Cardiol.* 2000; 35:1654-1660 (15.342)
9. Hamasaki S, Higano St, Suwaidi JA, Nishimura RA, Miyauchi K, Holmes DR Jr, Lerman A: Cholesterol-lowering treatment is associated with improvement in coronary vascular remodeling and endothelial function in patients with normal or mildly diseased coronary arteries. *Arterioacler thromb Vasc Biol.* 2000; 20: 737-743 (5.533)
10. Daida H, Kuwabara Y, Yokoi H, Nishikawa H, Takatsu F, Nakata Y, Kutsumi Y, Oshima S, Nishiyama S, Ishiwata S, Kato K, Nishimura S, Miyauchi K, Kanoh T, Yamaguchi H, and the PART Investigators: Effect of Probucol on repeat revascularization rate after percutaneous transluminal coronary angioplasty (from the Probucol Angioplasty Restenosis Trial [PART]). *Am J Cardiol.* 2000; 86: 550-552 (3.425)

11. Shimada K, Daida H, Mokuno H, Watanabe Y, Sawano M, Iwama Y, Seki E, Kurata T, Sato H, Ohashi S, Suzuki H, Miyauchi K, Takaya J, Sakurai H, Yamaguchi H: Association of seropositivity for antibody to Chlamydia-specific lipopolysaccharide and coronary artery disease in Japanese men. *Jpn Circ J.* 2001; 65:182-187 (3.685)
12. Panetta CJ, Miyauchi K, Berry D, Simari RD, Holmes DR, Schwartz RS, Caplice NM: A tissue-engineered stent for cell-based vascular gene transfer. *Human Gene Therapy* 2002; 13: 433-441 (3.625)
13. Inoue K, Cynshi O, Kawabe Y, Nakamura M, Miyauchi K, Kimura T, Daida H, Hamakubo T, Yamaguchi H, Kodama T: Effect of BO-653 and probucol on c-Myc and PDGF-A messenger RNA of the iliac artery after balloon denudation in cholesterol-fed rabbits. *Atherosclerosis.* 2002; 161: 353-363 (3.972)
14. Shimada K, Miyauchi K, Mokuno H, Miyazaki T, Seki E, Watanabe Y, Iwama Y, Shigekiyo M, Matsumoto M, Okazaki S, Tanimoto K, Kawamura M, Suzuki H, Kurata T, Sato H, Daida H. Predictive value of the adipocyte-derived plasma protein adiponectin for restenosis after elective coronary stenting. *Jpn Heart J.* 2002; 43: 85-91 (1.127)
15. Ozeki T, Miyauchi K, Takebayashi Y. Effectiveness of the DICOM structured based cardiac catheterization report. *Ann science of Graduate School of Shizuoka university.* 2004; 26: 21-27
16. Shimada K, Mokuno H, Matsunaga E, Miyazaki T, Sumiyoshi K, Kume A, Miyauchi K, Daida H. Predictive value of circulating oxidized LDL for cardiac events in type 2 diabetic patients with coronary artery disease. *Diabetes Care.* 2004; 27:843-4 (8.570)
17. Shimada K, Miyauchi K, Mokuno H, Watanabe Y, Iwama Y, Shigekiyo M, Matsumoto M, Okazaki S, Tanimoto K, Kurata T, Sato H, Daida H. Promoter polymorphism in the CD14 gene and concentration of soluble CD14 in patients with in-stent restenosis after elective coronary stenting. *Int J Cardiol.* 2004; 94:87-92 (6.175)
18. Yokoyama T, Miyauchi K, Kurata T, Sato H, Daida H. Effect of probucol on neointimal thickening in a stent porcine restenosis model. *Jpn Heart J.* 2004; 45: 305-13 (1.127)
19. Yokoyama T, Miyauchi K, Kurata T, Satoh H, Daida H. Inhibitory efficacy of pitavastatin on the early inflammatory response and neointimal thickening in a porcine coronary after stenting. *Atherosclerosis.* 2004; 174: 253-9 (3.971)
20. Shimada K, Mokuno H, Matsunaga E, Miyazaki T, Sumiyoshi K, Miyauchi K, Daida H. Circulating oxidized low-density lipoprotein is an independent predictor for cardiac event in patients with coronary artery disease. *Atherosclerosis.* 2004; 174: 343-7 (3.971)
21. Shimada K, Miyauchi K, Daida H. Early intervention with atorvastatin modulates TH1/TH2 imbalance in patients with acute coronary syndrome: from bedside to bench. *Circulation.* 2004; 109: e213-4 (14.948)

22. Okazaki S, Yokoyama T, Miyauchi K, Shimada K, Kurata T, Sato H, Daida Hb. Early Statin Treatment in Patients With Acute Coronary Syndrome. Demonstration of the Beneficial Effect on Atherosclerotic Lesions by Serial Volumetric Intravascular Ultrasound Analysis During Half a Year After Coronary Event: The ESTABLISH Study. *Circulation* 2004; 110: 1061-1068 (14.948)
23. Nijaman W, Miyauchi K, Kasai T, Kurata T, Sato H, Ohta H, Okazaki S, Yokoyama K, Kojima T, Akimoto Y, Daida H. Impact of aspirin treatment on long-term outcome (over 10-years) after percutaneous coronary intervention. *Int Heart J* 2006; 47: 37-45 (1.127)
24. Kasai T, Miyauchi K, Yokoyama T, Aihara K, Daida H. Efficacy of peroxisome proliferative activated receptor (PPAR)-alpha ligands, fenofibrate, on intimal hyperplasia and constrictive remodeling after coronary angioplasty in porcine models. *Atherosclerosis*. 2006; 188: 274-80 (3.971)
25. Miyauchi K, Kimura T, Morimoto T, Nakagawa Y, Yamagishi M, Ozaki Y, Hiro T, Daida H, Matsuzaki M. Japan Assessment of Pitavastatin and Atorvastatin in Acute Coronary Syndrome (JAPAN-ACS) –Rationale and Design-. *Cir J* 2006; 70: 1624-1628 (3.685)
26. Kasai T, Miyauchi K, Kurata T, Sato H, Ohta H, Okazaki S, Yokoyama K, Kojima T, Akimoto Y, Daida H. Prognostic value of the metabolic syndrome on Long-term (more than 10 years) Outcomes in Patient Undergoing Percutaneous Coronary Intervention. *Circ J* 2006; 70: 1531-1537 (3.685)
27. Kawamori R, Daida H, Tanaka Y, Miyauchi K, Kitagawa American Heart Associationyashi D, Kishimoto J, Ikeda S, Imai Y, Yamazaki T. Amlodipine versus angiotensin II receptor blocker; control of blood pressure evaluation trial in diabetics (ADVANCED-J). *BMC Cardiovascular Disorders*. 2006; 6: 39-50 (1.500)
28. Tanaka A, Azuma K, Toyofuku Y, Kurokawa A, Otsuka A, Mita T, Hirose T, Fujitani Y, Miyauchi K, Daida H, Kawamori R, Watada H. Insulin and nateglinide reduce monocyte adhesion to endothelial cells in Goto-Kakizaki rats exhibiting repetitive blood glucose fluctuation. *Biochem Biophys Res Commun*. 2006; 350: 195-201 (2.281)
29. Inoue K, Sugiyama A, Reid PC, Ito Y, Miyauchi K, Mukai S, Sagara M, Miyamoto K, Satoh H, Kohno I, Kurata T, Ota H, Mantovani American Heart Associationmakubo T, Daida H, Kodama T. Establishment of a High Sensitivity Plasma Assay for Human Pentraxin3 as a Marker for Unstable Angina Pectoris. *Arterioscler. Thromb. Vasc. Biol*. 2007; 27: 161-167 (5.533)
30. Kasai T, Miyauchi K, Kurata T, Sato H, Ohta H, Okazaki S, Yokoyama K, Kojima T, Akimoto Y, Daida H. Long-term (11-year) statin therapy following Percutaneous Coronary Intervention improves clinical outcomes but not associated with increased malignancy. *Int J Cardiol*. 2007; 114: 210-217 (6.175)
31. Ito M, Hatta K, Miyakawa K, Miyauchi K, Arai H. Prolonged and Fluctuating Giant T-wave inversion after electroconvulsive therapy. *J ECT* 2007; 23: 194-197

32. Kubota N; Kasai T; Miyauchi K, Njaman W; Kajimoto K; Akimoto Y; Kojima T; Yokoyama K; Kurata T; Daida H. Therapy with statins and aspirin enhances long-term outcome of percutaneous coronary intervention. *Heart and Vessels* 2008; 23; 35-9 (2.109)
33. Nagai R, Izumi T, Kurabayashi M, Daida H, Tojo T, Hasegawa A, Miyauchi K, Hayashi D, Kohro T, Okada Y, Yamazaki T; JCADII Investigators. Rationale and design of a study to examine lower targets for low-density lipoprotein-cholesterol and blood pressure in coronary artery disease patients. *Circ J* 2008; 72: 515-20 (3.685)
34. Kasai T, Miyauchi K, Kurata T, Okazaki S, Kojimoto K, Kubota N, Daida H. Impact of metabolic syndrome among patients with and without diabetes mellitus on long-term outcomes after percutaneous coronary intervention. *Hypertension Res* 2008; 31: 235-41 (2.935)
35. Kasai T, Miyauchi K, Kurata T, Sato H, Okazaki S, Kojimoto K, Kubota N, Daida H. Influence of Diabetes on >10 year outcomes after percutaneous coronary intervention. *Heart and Vessels* 2008; 23, 149-54 (2.109)
36. Kasai T, Miyauchi K, Yokoyama T, Kajimoto K, Sumiyoshi K, Kubota N, Ikeda E, Daida H. Pioglitazone attenuates neointimal thickening via suppression of the early inflammatory response in a porcine coronary after stenting. *Atherosclerosis* 2008; 197: 612-9 (3.971)
37. Kasai T, Miyauchi K, Kubota N, Tamura H, Kojima T; Yokoyama K; Kurata T, Daida H. The relationship between the metabolic syndrome defined by various criteria and the extent of coronary artery disease. *Atherosclerosis* 2008; 31: 235-41 (3.974)
38. Miyauchi K, Kasai T, Yokoyama T, Aihara K, Kurata T, Kajimoto K, Okazaki S, Ishiyama H, Daida H. Effectiveness of Statin Eluting-Stent on Early Inflammatory Response and Neointimal Thickness in a Porcine Coronary Model. *Circ J* 2008; 72: 832-8 (3.685)
39. Ikeda E, Kasai T, Kajimoto K, Miyauchi K, Kurata T, Kubota N, Kurata T, Amano A, Daida H. Dipyridamole therapy improves long-term survival after complete revascularization in patients with impaired cardiac function. –A propensity analysis-. *Circ J* 2008; 72: 1481-86 (3.685)
40. Kajimoto K, Miyauchi K, Kasai T, Hirose H, Yanagisawa N, Yamamoto T, Takazawa K, Niinami H, Hosoda Y, Daida H, Amano A. Metabolic Syndrome predicts 10-year mortality in non-diabetic patients following coronary artery bypass surgery. *Circ J* 2008; 72: 1481-86 (3.685)
41. Kajimoto K, Miyauchi K, Kasai T, Yanagisawa N, Yamamoto T, Kikuchi K, Nakatomi T, Iwamura H, Daida H, Amano A. Metabolic Syndrome is an Independent Risk Factor for Stroke and Acute Renal Failure after Coronary Artery Bypass Grafting. *J Thorac Cardiovasc Sur* 2009; 137: 658-663 (3.991)
42. Kasai T, Kajimoto K, Miyauchi K, Kurata T, Kubota N, Kurata T, Amano A, Daida H. Propensity analysis of 12 years outcome after bypass graft or balloon angioplasty in patients with multivessel coronary artery disease. *J Cardiol* 2008; 52: 186-194 (2.566)
43. Matsunaga E, Takaya N, Yokoyama T, Akimoto Y, Miyauchi K, Daida H. The Relationship

Between Coronary Artery Wall Thickness Measured by 64-slice Computed Tomography and Cardiovascular Risk Factors. *Circ J.* 2008; 73: 681-685 (3.685)

44. Miyauchi K, Daida H, Takaya N, Kawamori R, Hirose T, Ikeda F, Arai H, Ohishi H, Yoshida K, Yamamoto M, Hattori N, Urabe T, Maehara T, Suzuki M, Sase K, Hatsukami T, Yuan C. Rationale and design of the carotid plaque in human for all evaluation with aggressive rosvastatin therapy. *Circ J.* 2009; 73: 681-685 (3.685)

45. Kajimoto K, Miyauchi K, Kasai T, Kazunori Shimada, Yuko Kojima, Akie Shimada, Hiroshi Niinami, Atsushi Amano, Hiroyuki Daida First Human Experiments: Aggressive Statin Therapy Reduced Key Inflammatory Factors including c-Jun N-terminal kinase (JNK) and Dendritic Cell and Matrix Metalloproteinase Expression in Human Abdominal Aortic Aneurysmal Wall. *Atherosclerosis.* 2009; 206: 505-11 (3.974)

46. Kasai T, Miyauchi K, Kajimoto K, Kubota N, Yanagisawa N, Amano A, Daida H. Relationship between the metabolic syndrome and the incidence of stroke after complete coronary revascularization over a 10-year follow-up period. *Atherosclerosis* 2009; 207: 195-9 (3.974)

47. Tanaka Y, Daida H, Imai Y, Miyauchi K, Sato Y, Hiwatari M, Kitagawa A, Kishimoto J, Yamazaki T, Kawamori R. Morning home blood pressure may be a significant marker of nephropathy in Japanese patients with type 2 diabetes: ADVANCED-J study. *Hypertens Res* 2009; 32: 770-4 (2.935)

48. Dohi T, Miyauchi K, Kasai T, Kajimoto K, Kubota N, Tamura H, Yokoyama T, Kojima T, Yokoyama K, Kurata T, Daida H. Impact of Metabolic Syndrome on 10-Year Clinical Outcomes Among Patients With Acute Coronary Syndrome. *Circ J* 2009; 73: 1454-8 (3.685)

49. Kasai T, Miyauchi K, Kajimoto K, Kubota N, Dohi T, Kurata T, Amano A, Daida H. The adverse prognostic significance of the metabolic syndrome with and without hypertension in patients who underwent complete coronary revascularization. *J Hypertens* 2009; 27: 1017-24 (4.222)

50. Hiro T, Kimura T, Morimoto T, Miyauchi K, Nakagawa Y, Yamagishi M, Ozaki Y, Kazuo Kimura K, Saito S, Yamaguchi T, Daida H, Matsuzaki M and JAPAN-ACS Investigators Effect of Intensive Statin Therapy on Regression of Coronary Atherosclerosis in Patients With Acute Coronary Syndrome: A Multicenter Randomized Trial Evaluated by Volumetric Intravascular Ultrasound Using Pitavastatin Versus Atorvastatin (JAPAN-ACS [Japan Assessment of Pitavastatin and Atorvastatin in Acute Coronary Syndrome] Study) *J Am Coll Cardiol* 2009; 54: 293-302 (15.343)

51. Kasai T, Miyauchi K, Kajimoto K, Kubota N, Kurata T, Amano A, Daida H. The Impact of Pravastatin Therapy on Long-Term Outcome in Patients with Metabolic Syndrome Undergoing Complete Coronary Revascularization. *Circ J* 2009; 73: 2104-9 3.685)

52. Miyauchi K, Kojima T, Kawamura M, Suwa S, Okazaki S, Inoue K, Sumiyoshi M, Tanimoto K, Nakazato Y, Yamagami S, Daida H. Azelnidipine and Amlodipine Anti-Coronary

Atherosclerosis Trial in Hypertensive Patients Undergoing Coronary Intervention by Serial Volumetric Intravascular Ultrasound Analysis in Juntendo Medical University (ALPS-J) Cardiovasc Drugs Ther 2009; 23: 409-13 (2.952)

53. Heart Care Network Groups. Current trends in lifestyle-related disease management by general practitioners: a report from the "Heart Care Network" groups. J Atheroscler Thromb 2009; 16:799-806 (2.770)

54. Aihara K, Miyauchi K, Kasai T, Kubota N, Kajimoto K, Tamura H, Kojima T, Yokoyama K, Takeshi Kurata T, Amano A, Daida H. Long-term Efficacy of Pravastatin Therapy in Diabetic Patients Undergoing Complete Coronary Revascularization. J Atheroscler Thromb 2010; 17: 350-5 (2.770)

55. Dohi T, Miyauchi K, Okazaki S, Yokoyama T, Yanagisawa N, Tamura H, Kojima T, Yokoyama K, Kurata T, Daida H Early intensive statin treatment for six months improves long-term clinical outcomes in patients with acute coronary syndrome (Extended-ESTABLISH trial): a follow-up study. Atherosclerosis 2010; 210: 497-502 (3.974)

56. Yamagami S, Miyauchi K, Daida H. Impact of Initial Degree of Coronary Narrowing on Left Ventricular Function in First Myocardial Infarction. Int Heart J 2010; 51: 82-85 (1.127)

57. Miyauchi K, Morino Y, Tsukahara K, Origasa H, Daida H; PACIFIC steering committee members. The PACIFIC (Prevention of Atherothrombotic Incidents Following Ischemic Coronary attack) Registry: Rationale and design of a 2-year study in patients initially hospitalised with acute coronary syndrome in Japan. Cardiovasc Drugs Ther 2010; 24: 77-83 (2.952)

58. Hiro T, Kimura T, Morimoto T, Miyauchi K, Nakagawa Y, Yamagishi M, Ozaki Y, Kimura K, Saito S, Yamaguchi T, Daida H, Matsuzaki M; The JAPAN-ACS. Investigators Diabetes Mellitus Is a Major Negative Determinant of Coronary Plaque Regression During Statin Therapy in Patients With Acute Coronary Syndrome. Circ J 2010; 74: 1165-74 (3.685)

59. Miyazaki S, Kasai T, Miyauchi K, Miyazaki T, Akimoto Y, Takagi A, Aihara K, Kawamura M, Suwa S, Kojima S, Sumiyoshi M, Daida H. Changes of Matrix Metalloproteinase-9 Level Is Associated With Left Ventricular Remodeling Following Acute Myocardial Infarction Among Patients Treated With Trandolapril, Valsartan or Both. Circ J 2010; 74: 1158-64 (3.685)

60. Kubota N, Miyauchi K, Kasai T, Iesaki T, Kajimoto K, Sumiyoshi K, Dohi T, Daida H Synergistic effects of calcium-channel and Angiotensin-receptor blockers on endothelial function and inflammatory responses in a porcine drug-eluting stent model. Circ J 2010; 74: 1704-10 (3.685)

61. Ohashi T, Shibata R, Morimoto T, Kanashiro M, Ishii H, Ichimiya S, Hiro T, Miyauchi K, Nakagawa Y, Yamagishi M, Ozaki Y, Kimura T, Daida H, Murohara T, Matsuzaki M Correlation between circulating adiponectin levels and coronary plaque regression during aggressive lipid-lowering therapy in patients with acute coronary syndrome: subgroup analysis of JAPAN-ACS study. Atherosclerosis 2010; 212: 237-42 (3.974)

62. Arai H, Hiro T, Kimura T, Morimoto T, Miyauchi K, Nakagawa Y, Yamagishi M, Ozaki Y, Kimura K, Saito S, Yamaguchi T, Daida H, Matsuzaki M. More Intensive Lipid Lowering is Associated with Regression of Coronary Atherosclerosis in Diabetic Patients with Acute Coronary Syndrome. *J Atheroscler Thromb* 2010; 17: 1096-107 (2.770)
63. Miyazaki T, Shimada K, Miyauchi K, Kume A, Tanimoto K, Kiyanagi T, Sumiyoshi K, Hiki M, Mokuno H, Okazaki S, Sato H, Kurata T, Daida H. Effects of fenofibrate on lipid profiles, cholesterol ester transfer activity, and in-stent intimal hyperplasia in patients after elective coronary stenting. *Lipids Health Dis* 2010; 9: 122 (2.770)
64. Dohi T, Miyauchi K, Okazaki S, Yokoyama T, Yanagisawa N, Tamura H, Kojima T, Yokoyama K, Kurata T, Daida H. Higher Baseline LDL-C Levels Amplify the Short-Term Benefit of Early Intensive Statin Treatment in Acute Coronary Syndrome. *J Atheroscler Thromb* 2011; 18: 231-9 (2.770)
65. Kasai T, Miyauchi K, Kajimoto K, Kubota N, Dohi T, Tsuruta R, Ogita M, Yokoyama T, Amano A, Daida H. Prognostic significance of glomerular filtration rate estimated by the Japanese equation among patients who underwent complete coronary revascularization. *Hypertens Res* 2011; 34: 378-83(2.936)
66. Dohi T, Miyauchi K, Okazaki S, Yokoyama T, Yanagisawa N, Tamura H, Kojima T, Yokoyama K, Kurata T, Daida H. Plaque Regression Determined by Intravascular Ultrasound Predicts Long-Term Outcomes of Patients with Acute Coronary Syndrome. *J Atheroscler Thromb* 2011; 18: 231-9 (2.770)
67. Tsuruta R, Miyauchi K, Yamamoto T, Dohi S, Tambara K, Dohi T, Inaba H, Kuwaki K, Daida H, Amano A. Effect of preoperative hemoglobin A1c levels on long-term outcomes for diabetic patients after off-pump coronary artery bypass grafting. *J Cardiol* 2011; 57: 181-6 (2.566)
68. Masaki Y, Shimada K, Kojima T, Miyauchi K, Inoue K, Kiyanagi T, Hiki M, Fukao K, Hirose K, Ohsaka H, Kume A, Miyazaki T, Ohmura H, Ohsaka A, Daida H. J Clinical significance of the measurements of plasma N-terminal pro-B-type natriuretic peptide levels in patients with coronary artery disease who have undergone elective drug-eluting stent implantation. *J Cardiol* 2011; 57: 303-10 (2.566)
69. Dohi T, Miyauchi K, Iesaki T, Tsuruta R, Tsuboi S, Ogita M, Kubota N, Kasai T, Yokoyama T, Daida H. Candesartan With Pioglitazone Protects Against Endothelial Dysfunction and Inflammatory Responses in Porcine Coronary Arteries Implanted With Sirolimus-Eluting Stents. *Circ J* 2011; 75: 1098-1106 (3.685)
70. Dohi T, Miyauchi K, Okazaki S, Yokoyama T, Tamura H, Kojima T, Yokoyama K, Kurata T, Daida H. Long-term impact of mild chronic kidney disease in patients with acute coronary syndrome undergoing percutaneous coronary interventions. *Nephrol Dial Transplant* 2011; 26: 2906-11 (3.488)

71. Kojima T, Miyauchi K, Yokoyama T, Yokoyama K, Kurata T, Suwa S, Kawamura M, Tamura H, Okazaki S, Inoue K, Fujiwara Y, Sumiyoshi M, Tanimoto K, Nakazato Y, Yamagami S, Hiro T, Komiyama N, Daida H. Azelnidipine and Amlodipine Anti-Coronary Atherosclerosis Trial in Hypertensive Patients Undergoing Coronary Intervention by Serial Volumetric Intravascular Ultrasound Analysis in Juntendo University (ALPS-J). *Circ J* 2011; 75: 1071-79 (3.685)
72. Yokoyama K, Miyauchi K, Kawamura M, Kajimoto K, Dohi T, Yamagami S, Kano T, Amano A, Hosoda Y, Daida H String-sign in left internal thoracic artery is associated with regression in left main trunk stenosis after coronary artery bypass. *Int Heart J* 2011; 52: 84-7 (1.127)
73. Fukushima Y, Hirayama S, Ueno T, Dohi T, Miyazaki T, Ohmura H, Mokuno H, Miyauchi K, Miida T, Daida H Small dense LDL cholesterol is a robust therapeutic marker of statin treatment in patients with acute coronary syndrome and metabolic syndrome. *Clin Chim Acta* 2011; 412: 1423-7 (2.764)
74. Miyauchi K, Daida H. Intravascular Ultrasound Parameters in Non-Inferiority Trial. *Circ J* 2011; 75: 2283 (3.685)
75. Konishi H, Kasai T, Miyauchi K, Kajimoto K, Kubota N, Dohi T, Amano A, Daida H Association of Low Glomerular Filtration Rate With the Incidence of Stroke in Patients Following Complete Coronary Revascularization. *Circ J* 2011; 75: 2372-8 (3.685)
76. Sato H, Kasai T, Miyauchi K, Kubota N, Kajimoto K, Miyazaki T, Nishino A, Yaginuma K, Tamura H, Kojima T, Yokoyama K, Kurata T, Amano A, Daida H. Long-term outcomes of women with coronary artery disease following complete coronary revascularization. *J Cardiol* 2011; 58: 158-64 (2.566)
77. Kohro T, Yamazaki T, Izumi T, Daida H, Kurabayashi M, Miyauchi K, Tojo T, Nagai R and JCADII Investigator. Intensively Lowering Both Low-Density Lipoprotein Cholesterol and Blood Pressure Does Not Reduce Cardiovascular Risk in Japanese Coronary Artery Disease Patients. *Circ J* 2011; 75: 2062-70 (3.685)
78. Ogita M, Miyauchi K, Kurata T, Yokoyama K, Dohi T, Tuboi S, Miyazaki T, Nishino A, Yokoyama T, Daida H. Clinical Impact of Angiographic Restenosis After Bare-Metal Stent Implantation on Long-Term Outcomes in Patients With Coronary Artery Disease. *Circ J* 2011; 75: 2566-72 (3.685)
79. Hibi K, Kimura T, Kimura K, Morimoto T, Hiro T, Miyauchi K, Nakagawa Y, Yamagishi M, Ozaki Y, Saito S, Yamaguchi T, Daida H, Matsuzaki M; for the JAPAN-ACS Investigators. Clinically evident polyvascular disease and regression of coronary atherosclerosis after intensive statin therapy in patients with acute coronary syndrome: Serial intravascular ultrasound from the Japanese assessment of pitavastatin and atorvastatin in acute coronary syndrome (JAPAN-ACS) trial. *Atherosclerosis* 2011; 219: 743-9 (3.971)
80. Dohi T, Miyauchi K, Ohkawa R, Nakamura K, Kishimoto T, Miyazaki T, Nishino A,

Nakajima N, Yaginuma K, Tamura H, Kojima T, Yokoyama K, Kurata T, Shimada K, Yatomi Y, Daida H. Increased circulating plasma lysophosphatidic acid in patients with acute coronary syndrome. *Clin Chim Acta* 2012; 413: 610-6 (2.764)

81. Nishino A, Kasai T, Miyauchi K, Nakajima N, Yokoyama T, Yokoyama K, Kurata T, Daida H. Long-term effect of metabolic syndrome with and without diabetes mellitus on coronary revascularization in Japanese patients undergoing percutaneous coronary intervention. *Clin Cardiol* 2011; 34: 610-6 (2.225)

82. Dohi T, Miyauchi K, Okazaki S, Yokoyama T, Ohkawa R, Nakamura K, Yanagisawa N, Tsuboi S, Ogita M, Yokoyama K, Kurata T, Yatomi Y, Daida H. Decreased circulating lipoprotein-associated phospholipase A2 levels are associated with coronary plaque regression in patients with acute coronary syndrome. *Atherosclerosis* 2011; 219: 907-12 (3.971)

83. Kasai T, Miyauchi K, Kubota N, Kajimoto K, Amano A, Daida H. Probucol therapy improves long-term (>10-year) survival after complete revascularization: A propensity analysis. *Atherosclerosis* 2012; 220: 463-9 (3.971)

84. Yaginuma K, Kasai T, Miyauchi K, Kajimoto K, Amano A, Daida H. Propensity Score Analysis of 10-Year Long-term Outcome After Bypass Surgery or Plain Old Balloon Angioplasty in Patients With Metabolic Syndrome. *Int Heart J*. 2011; 52: 372-6 (1.127)

85. Ogita M, Miyauchi K, Dohi T, Wada H, Tsuboi S, Miyazaki T, Nishino A, Yokoyama T, Kojima T, Yokoyama K, Kurata T, Daida H. Gender-based outcomes among patients with diabetes mellitus after percutaneous coronary intervention in the drug-eluting stent era. *Int Heart J* 2011; 52: 348-52 (1.127)

86. Fukushima Y, Ohmura H, Mokuno H, Kajimoto K, Kasai T, Hirayama S, Miyauchi K, Miida T, Amano A, Daida H. Non-high-density lipoprotein cholesterol is a practical predictor of long-term cardiac death after coronary artery bypass grafting. *Atherosclerosis* 2012; 221: 206-11 (3.971)

87. Dohi S, Kajimoto K, Miyauchi K, Yamamoto T, Tambara K, Inaba H, Kuwaki K, Tamura H, Kojima T, Yokoyama K, Kurata T, Daida H, Amano A. Comparing outcomes after off-pump coronary artery bypass versus drug-eluting stent in diabetic patients. *J Cardiol* 2012; 59: 195-201 (2.566)

88. Kajimoto K, Miyauchi K, Yamamoto T, Daida H, Amano A. Meta-analysis of Randomized Controlled Trials on the Treatment of Unprotected Left Main Coronary Artery Disease: One-Year Outcomes with Coronary Artery Bypass Grafting Versus Percutaneous Coronary Artery Intervention with Drug-Eluting Stent. *J Card Surg* 2012; 27: 152-7 (0.888)

89. Miyauchi K, Daida H, Morimoto T, Hiro T, Kimura T, Nakagawa Y, Yamagishi M, Ozaki Y, Kadota K, Kimura K, Hirayama A, Kimura K, Hasegawa Y, Uchiyama S, Matsuzaki M; for the JAPAN-ACS Investigators. Reverse Vessel Remodeling But Not Coronary Plaque Regression Could

Predict Future Cardiovascular Events in ACS Patients With Intensive Statin Therapy. *Circ J* 2012; 76: 825-832 (3.685)

90. Nishizaki Y, Yamagami S, Haga K, Sesoko M, Yamashita H, Miyauchi K, Daida H. Usefulness of Prominently Projected Aortic Arch on Chest Radiograph to Predict Severe Tortuosity of the Right Subclavian or Brachiocephalic Artery in Patients Aged >44 Years Undergoing Coronary Angiography With a Right Radial Artery Approach. *Am J Cardiol* 2012; 110: 203-7 (3.425)

91. Matsumori R, Shimada K, Kiyonagi T, Hiki M, Fukao K, Hirose K, Ohsaka H, Miyazaki T, Kume A, Yamada A, Takagi A, Ohmura H, Miyauchi K, Daida H. Clinical significance of the measurements of urinary liver-type fatty acid binding protein levels in patients with acute coronary syndrome. *J Cardiol* 2012; 60: 168-73 (2.566)

92. Miyauchi K, Yamazaki T, Watada H, Tanaka Y, Kawamori R, Imai Y, Ikeda S, Kitagawa A, Ono Y, Murayama F, Choi JB, Suwa S, Hayashi D, Kishimoto J, Daida H; for the ADVANCED-J investigators. Management of Home Blood Pressure by Amlodipine Combined With Angiotensin II Receptor Blocker in Type 2 Diabetes. *Circ J* 2012; 76: 2159-66 (3.425)

93. Ohkawa R, Kishimoto T, Kurano M, Dohi T, Miyauchi K, Daida H, Nagasaki M, Uno K, Hayashi N, Sakai N, Matsuyama N, Nojiri T, Nakamura K, Okubo S, Yokota H, Ikeda H, Yatomi Y. Development of an enzymatic assay for sphingomyelin with rapid and automatable performances: Analysis in healthy subjects and coronary heart disease patients. *Clin Biochem* 2012; 16: 1463-70 (2.229)

94. Dohi T, Miyauchi K, Ohkawa R, Nakamura K, Thuboi S, Ogita M, Miyazaki T, Nishino A, Yokoyama K, Kurata T, Yatomi Y, Daida H. Higher lipoprotein-associated phospholipase A2 levels are associated with coronary atherosclerosis documented by coronary angiography. *Ann Clin Biochem* 2012; 413: 207-12 (2.080)

95. Takashima H, Ozaki Y, Morimoto T, Kimura T, Hiro T, Miyauchi K, Nakagawa Y, Yamagishi M, Daida H, Mizuno T, Asai K, Kuroda Y, Kosaka T, Kuhara Y, Kurita A, Maeda K, Amano T, Matsuzaki M; for the JAPAN-ACS Investigators. Clustering of Metabolic Syndrome Components Attenuates Coronary Plaque Regression During Intensive Statin Therapy in Patients With Acute Coronary Syndrome. Clustering of Metabolic Syndrome Components Attenuates Coronary Plaque Regression During Intensive Statin Therapy in Patients With Acute Coronary Syndrom. *Circ J* 2012; 76: 2840-7 (3.685)

96. Kasai T, Miyauchi K, Yanagisawa N, Kajimoto K, Kubota N, Ogita M, Tsuboi S, Amano A, Daida H. Mortality risk of triglyceride levels in patients with coronary artery disease. *Heart* 2012; 99: 22-9 (6.023)

97. Dohi T, Kasai T, Miyauchi K, Takasu K, Kajimoto K, Kubota N, Amano A, Daida H. Prognostic impact of chronic kidney disease on 10-year clinical outcomes among patients with acute coronary syndrome. *J Cardiol* 2012; 60: 438-42 (2.566)

98. Kimura T, Morimoto T, Natsuaki M, Shiomi H, Igarashi K, Kadota K, Tanabe K, Morino Y, Akasaka T, Takatsu Y, Nishikawa H, Yamamoto Y, Nakagawa Y, Hayashi Y, Iwabuchi M, Umeda H, Kawai K, Okada H, Kimura K, Simonton CA, Kozuma K; RESET Investigators. Comparison of everolimus-eluting and sirolimus-eluting coronary stents: 1-year outcomes from the Randomized Evaluation of Sirolimus-eluting Versus Everolimus-eluting stent Trial (RESET). *Circulation* 2012; 126: 1225-36 (14.948)
99. Miyauchi K, Lewis AD, Rajamannan NM, Kantor B, Camrud LJ, Camrud AR, Allen JL, Daida H, Yamaguchi H, David R, Holmes DR Jr, Schwartz RS, Cynshi O. EFFICACY OF LIPOPHILIC ANTIOXIDANT, BO-653, ON PROLIFERATION OF SMOOTH MUSCLE CELLS AND NEOINTIMAL THICKENING IN A PORCINE STENT MODEL. *Journal of Medical Research and Science* 2012; 2: 23-29
100. Tsuboi S, Miyauchi K, Kasai T, Ogita M, Dohi T, Miyazaki T, Yokoyama T, Kojima T, Yokoyama K, Kurata T, Daida H. Impact of Red Blood Cell Distribution Width on Long-Term Mortality in Diabetic Patients After Percutaneous Coronary Intervention. *Circ J* 2013; 77: 456-61 (3.685)
101. Ogita M, Miyauchi K, Dohi T, Tsuboi S, Miyazaki T, Yokoyama T, Yokoyama K, Shimada K, Kurata T, Jiang M, Bujo H, Daida H. Increased Circulating Soluble LR11 in Patients with Acute Coronary Syndrome. *Clin Chim Acta* 2013; 415: 191-4 (2.764)
102. Ogita M, Miyauchi K, Morimoto T, Daida H, Kimura T, Hiro T, Nakagawa Y, Yamagishi M, Ozaki Y, Matsuzaki M. Association between circulating matrix metalloproteinase levels and coronary plaque regression after acute coronary syndrome - Subanalysis of the JAPAN-ACS study. *Atherosclerosis* 2013; 226: 275-80 (3.971)
103. Fukushima Y, Daida H, Morimoto T, Kasai T, Miyauchi K, Yamagishi SI, Takeuchi M, Hiro T, Kimura T, Nakagawa Y, Yamagishi M, Ozaki Y, Matsuzaki M. Relationship between Advanced Glycation End Products and Plaque Progression in Patients with Acute Coronary Syndrome: The JAPAN-ACS Sub-study. *Cardiovasc Diabetol* 2013; 12: 1475
104. Daida H, Miyauchi K, Ogawa H, Yokoi H, Matsumoto M, Kitakaze M, Kimura T, Matsubara T, Ikari Y, Kimura K, Tsukahara K, Origasa H, Morino Y, Tsutsui H, Kobayashi M, Isshiki T; on behalf of the PACIFIC investigators Management and Two-Year Long-Term Clinical Outcome of Acute Coronary Syndrome in Japan. *Circ J* 2013; 77: 934-43
105. Ogita M, Miyauchi K, Miyazaki T, Naito R, Konishi H, Tsuboi S, Dohi T, Kasai T, Yokoyama T, Okazaki S, Kurata T, Daida H. Low high-density lipoprotein cholesterol is a residual risk factor associated with long-term clinical outcomes in diabetic patients with stable coronary artery disease who achieve optimal control of low-density lipoprotein cholesterol. *Heart Vessels* 2013; 29: 35-41
106. Dohi T, Miyauchi K, Ohkawa R, Nakamura K, Kurano M, Kishimoto T, Yanagisawa N,

Ogita M, Miyazaki T, Nishino A, Yaginuma K, Tamura H, Kojima T, Yokoyama K, Kurata T, Shimada K, Daida H, Yatomi Y. Increased lysophosphatidic acid levels in culprit coronary arteries of patients with acute coronary syndrome. *Atherosclerosis* 2013; 229: 192-7

107. Yang B, Daimon M, Ishii K, Kawata T, Miyazaki S, Hirose K, Ichikawa R, Chiang SJ, Suzuki H, Miyauchi K, Daida H. Prediction of Coronary Artery Stenosis at Rest in Patients With Normal Left Ventricular Wall Motion. *Int Heart J* 2013; 54: 266-272

108. Nishizaki Y, Daimon M, Miyazaki S, Suzuki H, Kawata T, Miyauchi K, Chiang SJ, Makinae H, Shinozaki T, Daida H. Clinical factors associated with classical symptoms of aortic valve stenosis. *J Heart Valve Dis* 2013; 22: 287-94

109. Chiang SJ, Daimon M, Ishii K, Kawata T, Miyazaki S, Hirose K, Ichikawa R, Miyauchi K, Yeh MH, Chang NC, Daida H. Assessment of Elevation of and Rapid Change in Left Ventricular Filling Pressure Using a Novel Global Strain Imaging Diastolic Index. *Circ J* 2013 Dec 10

110. Nishizaki Y, Shimada K, Tani S, Ogawa T, Ando J, Takahashi M, Yamamoto M, Shinozaki T, Miyauchi K, Nagao K, Hirayama A, Yoshimura M, Komuro I, Nagai R, Daida H. Significance of Imbalance in the Ratio of Serum n-3 to n-6 Polyunsaturated Fatty Acids in Patients With Acute Coronary Syndrome. *Am J Cardiol* 2014; 113: 441-5

111. Fujimoto S, Kondo T, Kodama T, Fujisawa Y, Groarke J, Kumamaru KK, Takamura K, Matsunaga E, Miyauchi K, Daida H, Rybicki FJ. A novel method for non-invasive plaque morphology analysis by coronary computed tomography angiography. *Int J Cardiovasc Imaging*. 2014; 30: 1373-82

112. Konishi H, Miyauchi K, Kasai T, Tsuboi S, Ogita M, Naito R, Katoh Y, Okai I, Tamura H, Okazaki S, Daida H. Long-term prognosis and clinical characteristics of young adults (≤ 40 years old) who underwent percutaneous coronary intervention. *J Cardiol*. 2014; 63:171-4

113. Chiang SJ, Daimon M, Ishii K, Kawata T, Miyazaki S, Hirose K, Ichikawa R, Miyauchi K, Yeh MH, Chang NC, Daida H. Assessment of elevation of and rapid change in left ventricular filling pressure using a novel global strain imaging diastolic index. *Circ J* 2014; 78 : 419-27

114. Nishizaki Y, Shimada K, Tani S, Ogawa T, Ando J, Takahashi M, Yamamoto M, Shinozaki T, Miyauchi K, Nagao K, Hirayama A, Yoshimura M, Komuro I, Nagai R, Daida H. Significance of imbalance in the ratio of serum n-3 to n-6 polyunsaturated fatty acids in patients with acute coronary syndrome. *Am J Cardiol*. 2014; 113: 441-5

115. Ueno Y, Yamashiro K, Tanaka Y, Watanabe M, Shimada Y, Kuroki T, Miyamoto N, Daimon M, Tanaka R, Miyauchi K, Daida H, Hattori N, Urabe T. Rationale and design of the EPISTEME trial: efficacy of post-stroke intensive rosuvastatin treatment for aortogenic embolic stroke. *Cardiovasc Drugs Ther* 2014; 28: 79-85

116. Naito R, Miyauchi K, Ogita M, Kasai T, Kawaguchi Y, Tsuboi S, Konishi H, Okazaki S, Kurata T, Daida H. Impact of admission glycemia and glycosylated hemoglobin A1c on long-term

clinical outcomes of non-diabetic patients with acute coronary syndrome. *J Cardiol.* 2014; 63: 106-11

117. Ogita M, Miyauchi K, Miyazaki T, Naito R, Konishi H, Tsuboi S, Dohi T, Kasai T, Yokoyama T, Okazaki S, Kurata T, Daida H. Low high-density lipoprotein cholesterol is a residual risk factor associated with long-term clinical outcomes in diabetic patients with stable coronary artery disease who achieve optimal control of low-density lipoprotein cholesterol. *Heart Vessels* 2014; 29: 35-41

118. Naito R, Miyauchi K, Konishi H, Tsuboi S, Ogita M, Kasai T, Tamura H, Okazaki S, Daida H. Comparison of long-term clinical outcomes between sirolimus- and paclitaxel-eluting stents in real-world clinical practice. *Heart Vessels.* 2014 Aug 13

119. Uchida T, Kasai T, Takagi A, Sekita G, Komiya K, Takeno K, Shigihara N, Shimada K, Miyauchi K, Fujitani Y, Daida H, Watada H. Prevalence of amiodarone-induced thyrotoxicosis and associated risk factors in Japanese patients. *Int J Endocrinol.* 2014; 2014: 534904

120. Dai Y, Konishi H, Takagi A, Miyauchi K, Daida H. Red cell distribution width predicts short- and long-term outcomes of acute congestive heart failure more effectively than hemoglobin. *Exp Ther Med* 2014; 8: 600-606

121. Shiomi H, Kozuma K, Morimoto T, Igarashi K, Kadota K, Tanabe K, Morino Y, Akasaka T, Abe M, Suwa S, Muramatsu T, Kobayashi M, Dai K, Nakao K, Uematsu M, Tarutani Y, Fujii K, Simonton CA, Kimura T; RESET Investigators. Long-term clinical outcomes after everolimus- and sirolimus-eluting coronary stent implantation: final 3-year follow-up of the randomized evaluation of sirolimus-eluting versus everolimus-eluting stent trial. *Circ Cardiovasc Interv* 2014; 7: 343-54

122. Nakajima N, Miyauchi K, Yokoyama T, Ogita M, Miyazaki T, Tamura H, Nishino A, Yokoyama K, Okazaki S, Kurata T, Suwa S, Daida H. Effect of combination of ezetimibe and a statin on coronary plaque regression in patients with acute coronary syndrome: ZEUS trial (eZEtimibe Ultrasound Study). *International Journal of Cardiology. Metabolic & ENDOCRINE* 2014; 3: 8-13

123. Ueno Y, Yamashiro K, Tanaka Y, Watanabe M, Miyamoto N, Shimada Y, Kuroki T, Tanaka R, Miyauchi K, Daida H, Hattori N, Urabe T. Rosuvastatin may stabilize atherosclerotic aortic plaque: transesophageal echocardiographic study in the EPISTEME trial. *Atherosclerosis.* 2015; 239: 476-82.

124. Konishi H, Miyauchi K, Kasai T, Tsuboi S, Ogita M, Naito R, Sai E, Fukushima Y, Katoh Y, Okai I, Tamura H, Okazaki S, Daida H. Impact of Lipoprotein(a) as Residual Risk on Long-Term Outcomes in Patients After Percutaneous Coronary Intervention. *Am J Cardiol.* 2015; 115: 157-60

125. Ogita M, Miyauchi K, Jiang M, Kasai T, Tsuboi S, Naito R, Konishi H, Dohi T, Yokoyama T, Okazaki S, Shimada K, Bujo H, Daida H. Circulating soluble LR11, a novel marker of smooth muscle cell proliferation, is enhanced after coronary stenting in response to vascular injury.

Atherosclerosis 2014; 237: 374-8

126. Kurano M, Suzuki A, Inoue A, Tokuhara Y, Kano K, Matsumoto H, Igarashi K, Ohkawa R, Nakamura K, Dohi T, Miyauchi K, Daida H, Tsukamoto K, Ikeda H, Aoki J, Yatomi Y. Possible Involvement of Minor Lysophospholipids in the Increase in Plasma Lysophosphatidic Acid in Acute Coronary Syndrome. *Arterioscler Thromb Vasc Biol* 2014 Nov 25
127. Konishi H, Miyauchi K, Kasai T, Tsuboi S, Ogita M, Naito R, Nishizaki Y, Okai I, Tamura H, Okazaki S, Isoda K, Daida H. Long-term Effect of β -blocker in ST-segment Elevation Myocardial Infarction in Patients with Preserved Left Ventricular Systolic Function: A propensity analysis. *Heart and Vessels*, 2015 in press (2.109)
128. Naito R, Miyauchi K, Ogita M, Tsuboi S, Konishi H, Dohi T, Kasai T, Tamura H, Okazaki S, Isoda K, Daida H. Clinical outcomes following percutaneous coronary intervention before and after introduction of drug-eluting stent. *Cardiovasc Interv Ther*. 2015 Apr
129. Sai E, Shimada K, Miyauchi K, Masaki Y, Kojima T, Miyazaki T, Kurata T, Ogita M, Tsuboi S, Yoshihara T, Miyazaki T, Ohsaka A, Daida H. Increased cystatin C levels as a risk factor of cardiovascular events in patients with preserved estimated glomerular filtration rate after elective percutaneous coronary intervention with drug-eluting stents. *Heart Vessels*. 2015 Apr 12
130. Mizuno A, Miyauchi K, Nishizaki Y, Yamazoe M, Komatsu I, Asano T, Mitsuhashi H, Nishi Y, Niwa K, Daida H. Impact of the augmentation time ratio on direct measurement of central aortic pressure in the presence of coronary artery disease. *Hypertens Res*. 2015 Apr 9.
131. Chiang SJ, Daimon M, Miyazaki S, Kawata T, Morimoto-Ichikawa R, Maruyama M, Ohmura H, Miyauchi K, Lee SL, Daida H. When and how aortic stenosis is first diagnosed: A single-center observational study. *J Cardiol*. 2015 Nov 18. pii: S0914-5087
132. Naito R, Miyauchi K, Konishi H, Tsuboi S, Ogita M, Dohi T, Kasai T, Tamura H, Okazaki S, Isoda K, Daida H. Appropriate Level of Low-Density Lipoprotein Cholesterol for Secondary Prevention of Coronary Artery Disease. *J Atheroscler Thromb*. 2015.
133. Nishitani-Yokoyama M, Miyauchi K, Shimada K, Miyazaki T, Ogita M, Okazaki S, Shioya M, Koba S, Tsujita H, Kobayashi Y, Daida H. Effects of Phase II Comprehensive Cardiac Rehabilitation on Coronary Plaque Volume After Acute Coronary Syndrome. *Int Heart J*. 2015; 56: 597-604.
134. Naito R, Miyauchi K, Konishi H, Tsuboi S, Ogita M, Dohi T, Kasai T, Tamura H, Okazaki S, Isoda K, Daida H. Temporal Trends in Clinical Outcome After Percutaneous Coronary Intervention 1984-2010 - Report From the Juntendo PCI Registry. *Circ J*. 2015
135. Naito R, Miyauchi K, Konishi H, Tsuboi S, Okazaki S, Daida H. Early definite stent thrombosis with everolimus-eluting stents. *Clin Case Rep*. 2015; 3: 854-7.
136. Al Shahi H, Shimada K, Miyauchi K, Yoshihara T, Sai E, Shiozawa T, Naito R, Aikawa T, Ouchi S, Kadoguchi T, Miyazaki T, Daida H. Elevated Circulating Levels of Inflammatory

Markers in Patients with Acute Coronary Syndrome. *Int J Vasc Med.* 2015; 15: 805375.

137. Konishi H, Miyauchi K, Dohi T, Tsuboi S, Ogita M, Naito R, Kasai T, Tamura H, Okazaki S, Isoda K, Daida H. Impact of stent length on clinical outcomes of first-generation and new-generation drug-eluting stents. *Cardiovasc Interv Ther.* 2015 Oct 23.

138. Naito R, Miyauchi K, Konishi H, Tsuboi S, Ogita M, Dohi T, Kajimoto K, Kasai T, Tamura H, Okazaki S, Isoda KC, Yamamoto T, Amano A, Daida H. Comparison of mortality between coronary artery bypass grafting and percutaneous coronary intervention with drug-eluting stents in elderly with diabetes and multivessel coronary disease. *Heart Vessels.* 2015 Sep 28.

139. Morinaga H, Miyazaki T, Tsutsui M, Kasai T, Takagi A, Kajino K, Miyauchi K, Daida H. Ischemic Cardiomyopathy with a Rapid Progression from Thrombotic Thrombocytopenic Purpura. *Intern Med.* 2015; 54: 2351-4.

140. Fujimoto S, Kondo T, Takamura K, Baber U, Shinozaki T, Nishizaki Y, Kawaguchi Y, Matsumori R, Hiki M, Miyauchi K, Daida H, Hecht H, Stone GW, Narula J. Incremental prognostic value of coronary computed tomographic angiography high-risk plaque characteristics in newly symptomatic patients. *J Cardiol.* 2015 Sep 7. pii: S0914-5087(15)00241-5.

141. Fujimoto S, Kondo T, Kumamaru KK, Shinozaki T, Takamura K, Kawaguchi Y, Matsumori R, Hiki M, Miyauchi K, Daida H, Rybicki FJ. Prognostic Value of Coronary Computed Tomography (CT) Angiography and Coronary Artery Calcium Score Performed Before Revascularization. *J Am Heart Assoc.* 2015; 4: e002264.

142. Miura M, Yamasaki M, Uemura Y, Yoshikawa M, Miyauchi K, Tanaka H, Miyachi H, Yamashita J, Suzuki M, Yamamoto T, Nagao K, Komuro I, Takayama M. Effect of Statin Treatment and Low-Density Lipoprotein-Cholesterol on Short-Term Mortality in Acute Myocardial Infarction Patients Undergoing Primary Percutaneous Coronary Intervention - Multicenter Registry From Tokyo CCU Network Database. *Circ J.* 2015 Dec 8

143. Suzuki M, Sumiyoshi T, Miyachi H, Yamashita J, Yamasaki M, Miyauchi K, Yamamoto T, Nagao K, Tomoike H, Takayama M. Effect of Coronary Thrombectomy in Cardiogenic Shock Complicating ST-Segment Elevation Myocardial Infarction. *Am J Cardiol.* 2015; 115: 1649-54.

144. Shibayama K, Daimon M, Watanabe H, Kawata T, Miyazaki S, Morimoto-Ichikawa R, Maruyama M, Chiang SJ, Miyauchi K, Daida H. Significance of Coronary Artery Disease and Left Ventricular Afterload in Unoperated Asymptomatic Aortic Stenosis. *Circ J.* 2015 Dec 24

145. Ogita M, Miyauchi K, Kasai T, Tsuboi S, Wada H, Naito R, Konishi H, Dohi T, Tamura H, Okazaki S, Yanagisawa N, Shimada K, Suwa S, Jiang M, Bujo H, Daida H. Prognostic impact of circulating soluble LR11 on long-term clinical outcomes in patients with coronary artery disease. *Atherosclerosis.* 2015; 244: 216-221

146. Kurano M, Dohi T, Nojiri T, Kobayashi T, Hirowatari Y, Inoue A, Kano K, Matsumoto H, Igarashi K, Nishikawa M, Miyauchi K, Daida H, Ikeda H, Aoki J, Yatomi Y. Blood levels of

serotonin are specifically correlated with plasma lysophosphatidylserine among the glycerol-lysophospholipids. *BBA Clin.* 2015 20; 4: 92-8.

147. Dai Y, Takagi A, Konishi H, Miyazaki T, Masuda H, Shimada K, Miyauchi K, Daida H. Long-term outcomes of rotational atherectomy in coronary bifurcation lesions. *Exp Ther Med.* 2015; 10: 2375-2383

148. Naito R, Miyauchi K, Konishi H, Tsuboi S, Ogita M, Dohi T, Kasai T, Tamura H, Okazaki S, Isoda K, Daida H. Gender difference in long-term clinical outcomes following percutaneous coronary intervention during 1984-2008. *Atherosclerosis.* 2015; 247: 105-110.

149. Naito R, Miyauchi K, Shitara J, Endo H, Wada H, Doi S, Konishi H, Tsuboi S, Ogita M, Dohi T, Kasai T, Tamura H, Okazaki S, Isoda K, Daida H. Temporal Trends in Clinical Outcomes Following Percutaneous Coronary Intervention in Patients with Renal Insufficiency. *J Atheroscler Thromb.* 2016 Feb 12.

150. Konishi H, Miyauchi K, Tsuboi S, Ogita M, Naito R, Dohi T, Kasai T, Tamura H, Okazaki S, Isoda K, Daida H Plasma lipoprotein(a) predicts major cardiovascular events in patients with chronic kidney disease who undergo percutaneous coronary intervention. *Int J Cardiol.* 2016; 205: 50-3

151. Wada H, Ogita M, Miyauchi K, Shitara J, Endo H, Doi S, Naito R, Konishi H, Tsuboi S, Dohi T, Kasai T, Tamura H, Okazaki S, Isoda K, Suwa S, Daida H. Impact of gender difference on long-term outcomes of percutaneous coronary intervention for coronary artery disease in patients under statin treatment. *Heart Vessels.* 2016 Apr 23

152. Konishi H, Miyauchi K, Kasai T, Tsuboi S, Ogita M, Naito R, Dohi T, Tamura H, Okazaki S, Daida H. Adequate time in therapeutic INR range using triple antithrombotic therapy is not associated with long-term cardiovascular events and major bleeding complications after drug-eluting stent implantation. *J Cardiol.* 2016 Mar 25. pii: S0914-5087

153. Tamura H, Miyauchi K, Dohi T, Tsuboi S, Ogita M, Kasai T, Okai I, Katoh Y, Miyazaki T, Naito R, Konishi H, Yokoyama K, Okazaki S, Isoda K, Kurata T, Daida H. Comparison of Clinical and Angiographic Outcomes After Bare Metal Stents and Drug-Eluting Stents Following Rotational Atherectomy. *Int Heart J.* 2016; 57: 150-7

154. Naito R, Miyauchi K, Daida H, Morimoto T, Hiro T, Kimura T, Nakagawa Y, Yamagishi M, Ozaki Y, Matsuzaki M; JAPAN-ACS Investigators. Impact of Total Risk Management on Coronary Plaque Regression in Diabetic Patients with Acute Coronary Syndrome. *J Atheroscler Thromb.* 2016 Mar

155. Ogita M, Miyauchi K, Onishi A, Tsuboi S, Wada H, Konishi H, Naito R, Dohi T, Kasai T, Kojima Y, Schwartz RS, Daida H. Development of Accelerated Coronary Atherosclerosis Model Using Low Density Lipoprotein Receptor Knock-Out Swine with Balloon Injury. *PLoS One.* 2016 Sep 15;11(9):e0163055

156. Wada H, Ogita M, Miyauchi K, Suwa S, Yamano M, Daida H Case report: Fulminant myocarditis associated with overwhelming pneumococcal infection. *Int J Cardiol.* 2016 Aug 18;223:706-707. doi: 10.1016/j.ijcard.2016.08.282
157. Naito R, Miyauchi K, Konishi H, Tsuboi S, Ogita M, Dohi T, Tamura H, Kasai T, Okazaki S, Isoda K, Daida H. Combined Effect of Body Mass Index and Renal Function on Long-Term Clinical Outcomes in Elderly Adults After Percutaneous Coronary Intervention. *J Am Geriatr Soc.* 2016 Sep;64(9):e39-41. doi: 10.1111/jgs.14223. Epub 2016 Aug 26
158. Wada H, Ogita M, Miyauchi K, Tsuboi S, Konishi H, Shitara J, Kunimoto M, Sonoda T, Iso T, Ebina H, Aoki E, Kitamura K, Tamura H, Suwa S, Daida H. Contemporary sex differences among patients with acute coronary syndrome treated by emergency percutaneous coronary intervention. *Cardiovasc Interv Ther.* 2016 Aug 8.
159. Kiso S, Naito R, Fukao K, Hiki M, Miyazaki T, Takagi A, Miyauchi K, Daida H. Rapidly progressed aortic stenosis in a patient with previous diagnosis of polycythemia vera and post-polycythemia vera myelofibrosis. *Clin Case Rep.* 2016 May 17;4(6):589-92. doi: 10.1002/ccr3.568
160. Nishizaki Y, Shimada K, Tani S, Ogawa T, Ando J, Takahashi M, Yamamoto M, Shinozaki T, Miyazaki T, Miyauchi K, Nagao K, Hirayama A, Yoshimura M, Komuro I, Nagai R, Daida H. Association between the docosahexaenoic acid to arachidonic acid ratio and acute coronary syndrome: a multicenter observational study. *BMC Cardiovasc Disord.* 2016 Jul 7;16(1):143. doi: 10.1186/s12874-016-0210-1
161. Noguchi K, Sakakibara M, Asakawa N, Tokuda Y, Kamiya K, Yoshitani T, Oba K, Miyauchi K, Nishizaki Y, Ogawa H, Yokoi H, Matsumoto M, Kitakaze M, Kimura T, Matsubara T, Ikari Y, Kimura K, Origasa H, Isshiki T, Morino Y, Daida H, Tsutsui H; PACIFIC investigators. Higher Hemoglobin A1c After Discharge Is an Independent Predictor of Adverse Outcomes in Patients With Acute Coronary Syndrome - Findings From the PACIFIC Registry. *Circ J.* 2016 Jun 24;80(7):1607-14. doi: 10.1253/circj.CJ-15-1126. Epub 2016 May 19
162. Takahashi S, Shimada K, Miyauchi K, Miyazaki T, Sai E, Ogita M, Tsuboi S, Tamura H, Okazaki S, Shiozawa T, Ouchi S, Aikawa T, Kadoguchi T, Al Shahi H, Yoshihara T, Hiki M, Isoda K, Daida H. Low and exacerbated levels of 1,5-anhydroglucitol are associated with cardiovascular events in patients after first-time elective percutaneous coronary intervention. *Cardiovasc Diabetol.* 2016 Oct 11;15(1):145
163. Konishi H, Miyauchi K, Shitara J, Endo H, Wada H, Doi S, Naito R, Tsuboi S, Ogita M, Dohi T, Kasai T, Okazaki S, Isoda K, Suwa S, Daida H. Impact of Lipoprotein(a) on Long-term Outcomes in Patients With Diabetes Mellitus Who Underwent Percutaneous Coronary Intervention. *Am J Cardiol.* 2016 Sep 13. pii: S0002-9149(16)31474-6. doi: 10.1016/j.amjcard.2016.08.012
164. Hassan A, Dohi T, Miyauchi K, Ogita M, Kurano M, Ohkawa R, Nakamura K, Tamura H,

- Isoda K, Okazaki S, Yatomi Y, Daida H Prognostic impact of homocysteine levels and homocysteine thiolactonase activity on long-term clinical outcomes in patients undergoing percutaneous coronary intervention. *J Cardiol.* 2016 Sep 27. pii: S0914-5087(16)30207-6. doi: 10.1016/j.jjcc.2016.08.013
165. Kuramitsu S, Miyauchi K, Yokoi H, Suwa S, Nishizaki Y, Yokoyama T, Nojiri S, Iwabuchi M, Shirai S, Ando K, Okazaki S, Tamura H, Watada H, Daida H Effect of sitagliptin on plaque changes in coronary artery following acute coronary syndrome in diabetic patients: The ESPECIAL-ACS study. *J Cardiol.* 2016 Sep 15. pii: S0914-5087(16)30194-0. doi: 10.1016/j.jjcc.2016.08.011
166. Shiomi H, Morimoto T, Kitaguchi S, Nakagawa Y, Ishii K, Haruna Y, Takamisawa I, Motooka M, Nakao K, Matsuda S, Mimoto S, Aoyama Y, Takeda T, Murata K, Akao M, Inada T, Eizawa H, Hyakuna E, Awano K, Shirotani M, Furukawa Y, Kadota K, Miyauchi K, Tanaka M, Noguchi Y, Nakamura S, Yasuda S, Miyazaki S, Daida H, Kimura K, Ikari Y, Hirayama H, Sumiyoshi T, Kimura T; ReACT Investigators. The ReACT Trial: Randomized Evaluation of Routine Follow-up Coronary Angiography After Percutaneous Coronary Intervention Trial. *JACC Cardiovasc Interv.* 2017 Jan 23;10(2):109-117. doi: 10.1016/j.jcin.2016.10.018. Epub 2016 Dec 28
167. Wada H, Dohi T, Miyauchi K, Shitara J, Endo H, Doi S, Naito R, Konishi H, Tsuboi S, Ogita M, Kasai T, Okazaki S, Isoda K, Suwa S, Daida H Impact of serum albumin levels on long-term outcomes in patients undergoing percutaneous coronary intervention *Heart Vessels.* 2017 Apr 20. doi: 10.1007/s00380-017-0981-8. [Epub ahead of print]
168. Wada H, Dohi T, Miyauchi K, Doi S, Naito R, Konishi H, Tsuboi S, Ogita M, Kasai T, Okazaki S, Isoda K, Suwa S, Daida H Independent and Combined Effects of Serum Albumin and C-Reactive Protein on Long-Term Outcomes of Patients Undergoing Percutaneous Coronary Intervention *Circ J.* 2017 Apr 19. doi: 10.1253/circj.CJ-17-0124.
169. Wada H, Dohi T, Miyauchi K, Doi S, Naito R, Konishi H, Tsuboi S, Ogita M, Kasai T, Hassan A, Okazaki S, Isoda K, Suwa S, Daida H. Prognostic Impact of the Geriatric Nutritional Risk Index on Long-Term Outcomes in Patients Who Underwent Percutaneous Coronary Intervention. *Am J Cardiol.* 2017 Mar 16. pii: S0002-9149(17)30291-6. PMID: 28388993
170. Hiki M, Miyazaki T, Shimada K, Sugita Y, Shimizu M, Aikawa T, Ouchi S, Shiozawa T, Takasu K, Takahashi S, Takagi A, Miyauchi K, Daida H. Significance of Serum Polyunsaturated Fatty Acid Level Imbalance in Patients with Acute Venous Thromboembolism. *J Atheroscler Thromb.* 2017 Mar 24. doi: 10.5551/jat.37424. [Epub ahead of print]
171. Suwa S, Ogita M, Miyauchi K, Sonoda T, Konishi H, Tsuboi S, Wada H, Naito R, Dohi T, Kasai T, Okazaki S, Isoda K, Daida H Impact of Lipoprotein (a) on Long-Term Outcomes in Patients with Coronary Artery Disease Treated with Statin After a First Percutaneous Coronary Intervention *J Atheroscler Thromb.* 2017 Mar 17. doi: 10.5551/jat.38794